



A/Box Seq
PATENT

Docket No. GM10025



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Nicola G. Wallis, et al

For: MurC

Assistant Commissioner for Patents

Box Patent Application

Washington, D.C. 20231

COVER SHEET FOR FILING PATENT APPLICATION

Enclosed herewith for filing in the United States Patent and Trademark Office are the Specification and Declaration and Power of Attorney for the patent application identified below:

1. The names of the inventors are: Nicola G. Wallis and Martin K. R. Burnham
2. The addresses of the inventors are:

Nicola G. Wallis
219 Sugartown Road, K203
Wayne, PA 19087; USA
(Citizen of the United Kingdom)

Martin K. R. Burnham
2927 Tanglewood Lane
Norristown, PA 19403; USA
(Citizen of the United Kingdom)
3. The title of the invention is: MurC
4. The name, registration and telephone number of the attorney is:
Attorney: Arthur E. Jackson
Registration No. 34,354
Telephone No. 609-520-3254

09103287-062398



5. The Docket Number used to identify this application is: GM10025
6. The correspondence address for this application is:
Stephen T. Falk
Dechert, Price & Rhoads
4000 Bell Atlantic Tower, 1717 Arch Street
Philadelphia, PA 19103-2793

This invention was not made by an agency of the United States Government, or under contract with an agency of the United States Government.

This application claims benefit to US Provisional Patent Application Number 60/052,720, filed July 3, 1997.

There are 73 pages of specification, 0 sheets of drawings, 17 claims and a CRF of the gene sequence listing accompanying this cover sheet. The CRF is identical to the paper copy of the sequence listing and contains no new matter.

Stephen T. Falk
Registration No. 36,795
Dechert, Price & Rhoads
4000 Bell Atlantic Tower
1717 Arch Street
Philadelphia, PA 19103-2793, USA
Phone: 215/994-2488

Date: June 23, 1998

CERTIFICATION UNDER 37 CFR 1.10

I hereby certify that this correspondence and the documents referred to as attached therein are being deposited with the United States Postal Service on **June 23, 1998**, in an envelope as "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 CFR 1.10, Mailing Label Number **EE 472 817 877 US** addressed to: Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231.

Robert Ballarini

Docket No. GM10025

INVENTORS:

**Nicola G. Wallis
219 Sugartown Road, K203
Wayne, PA 19087; USA
(Citizen of the United Kingdom)**

**Martin K. R. Burnham
2927 Tanglewood Lane
Norristown, PA 19403; USA
(Citizen of the United Kingdom)**

TITLE: MurC

MurC

RELATED APPLICATIONS

5

This application claims benefit to US Provisional Patent Application Number 60/052,720, filed July 3, 1997.

FIELD OF THE INVENTION

10

15

This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, the invention relates to polynucleotides and polypeptides of the MurC family, as well as their variants, hereinafter referred to as "MurC," "MurC polynucleotide(s)," and "MurC polypeptide(s)" as the case may be.

BACKGROUND OF THE INVENTION

20

25

It is particularly preferred to employ Staphylococcal genes and gene products as targets for the development of antibiotics. The Staphylococci make up a medically important genera of microbes. They are known to produce two types of disease, invasive and toxigenic. Invasive infections are characterized generally by abscess formation effecting both skin surfaces and deep tissues. *S. aureus* is the second leading cause of bacteremia in cancer patients. Osteomyelitis, septic arthritis, septic thrombophlebitis and acute bacterial endocarditis are also relatively common. There are at least three clinical conditions resulting from the toxigenic properties of Staphylococci. The manifestation of these diseases result from the actions of exotoxins as opposed to tissue invasion and bacteremia. These conditions include: Staphylococcal food poisoning, scalded skin syndrome and toxic shock syndrome.

30

The frequency of *Staphylococcus aureus* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Staphylococcus aureus* strains which are resistant to some or all of the standard antibiotics.

This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

Moreover, the drug discovery process is currently undergoing a fundamental revolution as it embraces "functional genomics," that is, high throughput genome- or gene-based biology. This approach is rapidly superseding earlier approaches based on "positional cloning" and other methods. Functional genomics relies heavily on the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available as well as from other sources. There is a continuing and significant need to identify and characterize further genes and other polynucleotides sequences and their related polypeptides, as targets for drug discovery.

Clearly, there exists a need for polynucleotides and polypeptides, such as the MurC embodiments of the invention, that have a present benefit of, among other things, being useful to screen compounds for antimicrobial activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists to find ways to prevent, ameliorate or correct such infection, dysfunction and disease.

The enzyme UDP-N-acetylmuramate:L-alanine ligase, encoded by the gene MurC catalyses the addition of the first amino acid (L-alanine) of the peptide moiety in peptidoglycan biosynthesis. L-alanine is added to UDP-N-acetyl muramate with the concomittant hydrolysis of ATP to yield UDP-N-acetylmuramyl-L-alanine, ADP and phosphate. The gene has been cloned and sequenced from *Escherichia coli* and the corresponding protein has been over-expressed, purified and kinetically characterised (Liger, D., Masson, A., Blanot, D., van Heijenoort, J. & Parquet, C., Eur. J. Biochem. 230, 80-87). The kinetic mechanism of this enzyme has been investigated. (Falk, P.J., Ervin, K.M., Volk, K.S. & Ho, H.T. (1996) Biochemistry, 35, 1417-1422). The gene sequences of MurC from *Bacillus subtilis* and *Haemophilus influenzae* are also known.

The discovery of a MurC homologue in the human pathogen *Staphylococcus aureus* will allow us to produce UDP-N-acetylmuramate:L-alanine ligase enzyme which can then be used to screen for novel antibiotics. Inhibitors of this protein have utility in anti-bacterial therapy as they will prevent the construction of the bacterial cell wall.

Certain of the polypeptides of the invention possess significant amino acid sequence homology to a known MurC protein.

SUMMARY OF THE INVENTION

The present invention relates to MurC, in particular MurC polypeptides and MurC polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including treatment of microbial diseases, amongst others. In a further aspect, the invention relates to methods for identifying agonists and antagonists using the materials provided by the invention, and for treating microbial infections and conditions associated with such infections with the identified agonist or antagonist compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such as assays for detecting MurC expression or activity.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The invention relates to MurC polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a MurC of *Staphylococcus aureus*, which is related by amino acid sequence homology to MurC polypeptide. The invention relates especially to MurC having the nucleotide and amino acid sequences set out in Table 1 as SEQ ID NO: 1 or 3 and SEQ ID NO: 2 or 4 respectively.

TABLE 1
MurC Polynucleotide and Polypeptide Sequences

(A) *Staphylococcus aureus* MurC polynucleotide sequence [SEQ ID NO:1].
5' -ATGAGTAAGGAGTTTATATAATGACACACTATCATTTTGTCTCGGAATTAAAGGTTCTGGC
ATGAGTTCATTAGCACAAATCATGCATGATTTAGGACATGAAGTTCAAGGATCGGATATT
GAGAACTACGTATTTACAGAAGTTGCTCTTAGAAATAAGGGGATAAAAATATTACCATT
GGTGCTAATAACATAAAAGAAGATATGGTAGTTATACAAGGTAATGCATTGCGAGTAGC

CATGAAGAAATAGTACGTGCACATCAATTGAAATTAGATGTTGTAAGTTATAATGATTTT
 TTAGGACAGATTATTGATCAATATACTTCAGTAGCTGTAAGTGGTGCACATGGTAAACT
 TCTACAACAGGTTTATTATCACATGTTATGAATGGTGATAAAAAAGACTTCATTTTAAATT
 GGTGATGGCACAGGTATGGGATTGCCTGAAAGTGATTATTTTCGCTTTTGAGGCATGTGAA
 5 TATAGACGTCACTTTTAAAGTTATAAACCTGATTACGCAATTATGACAAATATTGATTTT
 GATCATCCTGATTATTTCAAAGATATTAATGATGTTTTTGGATGCATTCCAAGAAATGGCA
 CATAATGTTAAAAAAGGTATTATTGCTTGGGGTGATGATGAACATCTACGTAAAATTGAA
 GCAGATGTTCCAATTTATTACTATGGATTTAAAGATTTCGGATGACATTTATGCTCAAAT
 ATTCAAATTACGGATAAAGGTACTGCTTTTGGATGTGTATGTGGATGGTGAGTTTATGAT
 10 CACTTCCTGTCTCCACAATATGGTGACCATACAGTTTTAAATGCATTAGCTGTAATTGCG
 ATTAGTTATTTAGAGAAGCTAGATGTTACAAATATTAAAGAAGCATTAGAAACGTTTGGT
 GGTGTTAAACGTCGTTTCAATGAACTACAATTGCAAATCAAGTTATTGTAGATGATTAT
 GCACACCATCCAAGAGAAATTAGTGCTACAATTGACACAGCACGAAAGAAATATCCACAT
 AAAGAAGTTGTTGCAGTATTTCAACCACACACTTCTCTAGAACACAAGCATTTTAAAT
 15 GAATTTGCAGAAAGTTTATGTAAAGCAGATCGTGTATTCTTATGTGAAATTTTGGCTCA
 ATTAGAGAAAATTCTGGCGCATTAAACGATACAAGATTTAATTGATAAAATTGGAGGTGCA
 TCGTTCATTAATGAAGATCTTATTAATGTATTAGAACAATTTGATAATGCTGTTGTTT
 TTTATGGGTGCAGGTGATATTCAAAAATTACAAAATGCATATTTAGATAAATTAGGCATG
 AAAAAATGCGTTTTAATATGTTTATAATAGAG-3'

20

(B) *Staphylococcus aureus* MurC polypeptide sequence deduced from a polynucleotide sequence in this table [SEQ ID NO:2].

NH₂-MTHYHFVGKSGMSSLAQIMHDLGHEVQGSIDIENYVFTEVALRNKGKILPFGANNIKE
 DMVVIQGNFASSHEEIVRAHQLKLDVVSYNDFLGQIIDQYTSVAVTGAHGKTSTTGLLS
 25 HVMNGDKKTSFLIGDGTGMGLPESDYFAFEACEYRRHFLSYKPDYAIMTNIDFDHPDYFK
 DINDVFDAFQEMAHNVKKGI IAWGDDEHLRKIEADVPIYYYGFKDSDDIYAQNIQITDKG
 TAFDVYVDGEFYDHFSLSPQYGDHTVLNALAVIAISYLEKLDVTNIKEALETFGGVKRRFN
 ETTIANQVIVDDYAHHPREISATIDTARKKYPHKEVVAVFQPHTFSRQTQAFLEFAESLC
 KADRVFLCEIFGSIRENSGALTIQDLIDKIGGASFINEDLINVLEQFDNAVVLFMGAGDI
 30 QKLQNAYLDKLGKNAF-COOH

30

(C) *Staphylococcus aureus* MurC ORF sequence [SEQ ID NO:3].

5'-
 ATTTAAAGATTCGGATGACATTTATGCTCAAATATTTCAAATTACGGATAAAGGTACTGCTGTTGATGTGTA
 35 TGTGGATGGTGAGTTTTATGATCACTTCCTGTCTCCACAATATGGTGACCATACAGTTTTAAATGCATTAGC
 TGTAAATGCGATTAGTTATTTAGAGAAGCTAGATGTTACAAATATTAAAGAAGCATTAGAAACGTTTGGTGG
 TGTAAACGTCGTTTCAATGAACTACAATTGCAAATCAAGTTATTGTAGATGATTATGCACACCATCCAAG
 AGAAATTAGTGC

5 TACAATTGACACAGCACGAAAGAAATATCCACATAAAGAAGTTGTTGCAGTATTTCAACCACACACTTTCTC
TAGAACACAAGCATTTTTAAATGAATTTGCAGAAAGTTTAAGTAAAGCAGATCGTGTATTCTTATGTGAAAT
TTTTGGATCAATTAGAGAAAATACTGCCGCATTAACGATACAAGATTTAATTGATAAAATTGAAGGTGCATC
GTTAATTAATGAAGATTCTATTAATGTATTAGAACAATTTGATAATGCTGTTGTTTTATTTATGGGTGCAGG
TGATATTCAAAA
ATTACAAAATGCATATTTAGATAAATTAGGCATGAAAAATGCGTTTTAATATGTTTATAA-3'

10 (D) *Staphylococcus aureus* MurC polypeptide sequence deduced from a polynucleotide ORF
sequence in this table [SEQ ID NO:4].

NH₂-
FKDSDDIYAQIFQITDKGTAVDVYVDGEFYDHFSLSPQYGDHTVLNALAVIAISYLEKLDVTNIKEALETFGG
VKRRFNETTIANQVIVDDYAHHPREISATIDTARKKYPHKEVVAVFQPHTFSRTQAFLNEFAESLSKADRVF
LCEIFGSIRENTGALTIQDLIDKIEGASLINEDSINVLEQFDNAVVLFMGAGDIQKLQAYLDKLGKNAF-
15 COOH

Deposited materials

20 A deposit containing a *Staphylococcus aureus* WCUH 29 strain has been deposited with the
National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St. Machar
Drive, Aberdeen AB2 1RY, Scotland on 11 September 1995 and assigned NCIMB Deposit No.
40771, and referred to as *Staphylococcus aureus* WCUH29 on deposit. . The *Staphylococcus*
aureus strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited
strain."

25 The deposited strain contains the full length MurC gene. The sequence of the
polynucleotides contained in the deposited strain, as well as the amino acid sequence of any
polypeptide encoded thereby, are controlling in the event of any conflict with any description of
sequences herein.

30 The deposit of the deposited strain has been made under the terms of the Budapest Treaty
on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent
Procedure. The strain will be irrevocably and without restriction or condition released to the public
upon the issuance of a patent. The deposited strain is provided merely as convenience to those of
skill in the art and is not an admission that a deposit is required for enablement, such as that
required under 35 U.S.C. §112.

A license may be required to make, use or sell the deposited strain, and compounds derived therefrom, and no such license is hereby granted.

In one aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Staphylococcus aureus* WCUH 29 strain, which polypeptide is contained in the deposited strain. Further provided by the invention are MurC polynucleotide sequences in the deposited strain, such as DNA and RNA, and amino acid sequences encoded thereby. Also provided by the invention are MurC polypeptide and polynucleotide sequences isolated from the deposited strain.

Polypeptides

MurC polypeptide of the invention is substantially phylogenetically related to other proteins of the MurC family.

In one aspect of the invention there are provided polypeptides of *Staphylococcus aureus* referred to herein as "MurC" and "MurC polypeptides" as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

Among the particularly preferred embodiments of the invention are variants of MurC polypeptide encoded by naturally occurring alleles of the MurC gene.

The present invention further provides for an isolated polypeptide which:

- (a) comprises or consists of an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97, 98, 99 or 99.5% or exact identity, to that of SEQ ID NO:2 over the entire length of SEQ ID NO:2;
- (b) a polypeptide encoded by an isolated polynucleotide comprising or consisting of a polynucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or exact identity to SEQ ID NO:1 over the entire length of SEQ ID NO:1, or the entire length of SEQ ID NO:1 encoding SEQ ID NO:2;
- (c) a polypeptide encoded by an isolated polynucleotide comprising or consisting of a polynucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity,

even more preferably at least 97, 98, 99 or 99.5% or exact identity, to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ ID NO:2; or

(d) a polypeptide encoded by an isolated polynucleotide comprising or consisting of a polynucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or exact identity, to SEQ ID NO:1 over the entire length of SEQ ID NO:3;


(e) a polypeptide encoded by an isolated polynucleotide comprising or consisting of a polynucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or exact identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3; or

(f) a polypeptide encoded by an isolated polynucleotide comprising or consisting of a polynucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or exact identity, to the amino acid sequence of SEQ ID NO:4, over the entire length of SEQ ID NO:4;

(g) comprises or consists of an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97, 98, 99 or 99.5% or exact identity, to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:4.

The polypeptides of the invention include a polypeptide of Table 1 [SEQ ID NO:2 or 4] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of MurC, and also those which have at least 70% identity to a polypeptide of Table 1 [SEQ ID NO:1 or 3] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and more preferably at least 90% identity to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and still more preferably at least 95% identity to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

5 The invention also includes a polypeptide consisting of or comprising a polypeptide of the formula:



10 wherein, at the amino terminus, X is hydrogen, a metal or any other moiety described herein for modified polypeptides, and at the carboxyl terminus, Y is hydrogen, a metal or any other moiety described herein for modified polypeptides, R₁ and R₃ are any amino acid residue or modified amino acid residue, m is an integer between 1 and 1000 or zero, n is an integer between 1 and 1000 or zero, and R₂ is an amino acid sequence of the invention, particularly an amino acid sequence selected from Table 1 or modified forms thereof. In the formula above, R₂ is oriented so that its amino terminal amino acid residue is at the left, covalently bound to R₁, and its carboxy terminal amino acid residue is at the right, covalently bound to R₃. Any stretch of amino acid residues denoted by either R₁ or R₃, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer. Other preferred embodiments of the invention are provided where m is an integer between 1 and 50, 100 or 500, and n is an integer between 1 and 50, 100, or 500.

15 It is most preferred that a polypeptide of the invention is derived from *Staphylococcus aureus*, however, it may preferably be obtained from other organisms of the same taxonomic genus. A polypeptide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

20 A fragment is a variant polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with MurC polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

25 Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NO:2 or 4], or of variants thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence. Degradation forms of the polypeptides of the invention produced by or in a host cell, particularly a *Staphylococcus aureus*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and

30

coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Further preferred fragments include an isolated polypeptide comprising an amino acid
5 sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO: 2, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO: 2.

Also preferred are biologically active fragments which are those fragments that mediate
10 activities of MurC, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of *Staphylococcus aureus* or the ability to initiate, or maintain cause Disease in an individual, particularly a human.

15 Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

In addition to the standard single and triple letter representations for amino acids, the term "X" or "Xaa" may also be used in describing certain polypeptides of the invention. "X" and
20 "Xaa" mean that any of the twenty naturally occurring amino acids may appear at such a designated position in the polypeptide sequence.

Polynucleotides

It is an object of the invention to provide polynucleotides that encode MurC polypeptides, particularly polynucleotides that encode the polypeptide herein designated MurC.

25 In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding MurC polypeptides comprising a sequence set out in Table 1 [SEQ ID NO:1 or 3] which includes a full length gene, or a variant thereof. The Applicants believe that this full length gene is essential to the growth and/or survival of an organism which possesses it, such as *Staphylococcus aureus*.

30 As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing MurC polypeptides and polynucleotides, particularly

5 *Staphylococcus aureus* MurC polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B- and Z-DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

Another aspect of the invention relates to isolated polynucleotides, including at least one full length gene, that encodes a MurC polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2 or 4] and polynucleotides closely related thereto and variants thereof.

10 In another particularly preferred embodiment of the invention there is a MurC polypeptide from *Staphylococcus aureus* comprising or consisting of an amino acid sequence of Table 1 [SEQ ID NO:2 or 4], or a variant thereof.

Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NO:1 or 3], a polynucleotide of the invention encoding MurC polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Staphylococcus aureus* WCUH 29 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in Table 1 [SEQ ID NO:1 or 3], typically a library of clones of chromosomal DNA of *Staphylococcus aureus* WCUH 29 in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA sequencing may also be performed to obtain a full length gene sequence.

15
20
25
30

Illustrative of the invention, each polynucleotide set out in Table 1 [SEQ ID NO:1 or 3] was discovered in a DNA library derived from *Staphylococcus aureus* WCUH 29.

Moreover, each DNA sequence set out in Table 1 [SEQ ID NO:1 or 3] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID NO:2 or 4] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known to those skilled in the art. The polynucleotide of SEQ ID NO: 1, between nucleotide number 22 and the stop codon which begins at nucleotide number 1333 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:

- (a) a polynucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or exact identity to SEQ ID NO:1 over the entire length of SEQ ID NO:1, or the entire length of SEQ ID NO:1 encoding SEQ ID NO:2;
- (b) a polynucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or 100% exact, to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ ID NO:2; or
- (c) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or 100% identity, to SEQ ID NO:1 over the entire length of SEQ ID NO:3;
- (d) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or exact identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3; or
- (e) a polynucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97, 98, 99 or 99.5% or exact identity, to the amino acid sequence of SEQ ID NO:4, over the entire length of SEQ ID NO:4.

5 A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than *Staphylococcus aureus*, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled or detectable probe consisting of or comprising the sequence of SEQ ID NO: 1 or 3 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

10 The invention provides a polynucleotide sequence identical over its entire length to a coding sequence (open reading frame) in Table 1 [SEQ ID NO:1 or 3]. Also provided by the invention is a coding sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence. The polynucleotide of the invention may also contain at least one non-coding sequence, including for example, but not limited to at least one non-coding 5' and 3' sequence, such as the transcribed but non-translated sequences, termination signals (such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson *et al.*, *Cell* 37: 767 (1984), both of which may be useful in purifying polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

25 A preferred embodiment of the invention is a polynucleotide of consisting of or comprising nucleotide 22 to the nucleotide immediately upstream of or including nucleotide 1333 set forth in SEQ ID NO:1 of Table 1, both of which encode the MurC polypeptide.

The invention also includes a polynucleotide consisting of or comprising a polynucleotide of the formula:

30
$$X-(R_1)_m-(R_2)-(R_3)_n-Y$$

wherein, at the 5' end of the molecule, X is hydrogen, a metal or a modified nucleotide residue, or together with Y defines a covalent bond, and at the 3' end of the molecule, Y is hydrogen, a metal, or a modified nucleotide residue, or together with X defines the covalent bond, each occurrence of R₁ and R₃ is independently any nucleic acid residue or modified nucleic acid residue, m is an integer between 1 and 3000 or zero, n is an integer between 1 and 3000 or zero, and R₂ is a nucleic acid sequence or modified nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from Table 1 or a modified nucleic acid sequence thereof. In the polynucleotide formula above, R₂ is oriented so that its 5' end nucleic acid residue is at the left, bound to R₁, and its 3' end nucleic acid residue is at the right, bound to R₃. Any stretch of nucleic acid residues denoted by either R₁ and/or R₂, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer. Where, in a preferred embodiment, X and Y together define a covalent bond, the polynucleotide of the above formula is a closed, circular polynucleotide, which can be a double-stranded polynucleotide wherein the formula shows a first strand to which the second strand is complementary. In another preferred embodiment m and/or n is an integer between 1 and 1000. Other preferred embodiments of the invention are provided where m is an integer between 1 and 50, 100 or 500, and n is an integer between 1 and 50, 100, or 500.

It is most preferred that a polynucleotide of the invention is derived from *Staphylococcus aureus*, however, it may preferably be obtained from other organisms of the same taxonomic genus.

A polynucleotide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Staphylococcus aureus* MurC having an amino acid sequence set out in Table 1 [SEQ ID NO:2 or 4]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode variants of a polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2 or 4]. Fragments of a polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

5 Further particularly preferred embodiments are polynucleotides encoding MurC variants, that have the amino acid sequence of MurC polypeptide of Table 1 [SEQ ID NO:2 or 4] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of MurC polypeptide.

10 Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding MurC polypeptide having an amino acid sequence set out in Table 1 [SEQ ID NO:2 or 4], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding MurC
15 polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least
20 99% being the more preferred.

Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of Table 1 [SEQ ID NO:1 or 3].

In accordance with certain preferred embodiments of this invention there are provided
25 polynucleotides that hybridize, particularly under stringent conditions, to MurC polynucleotide sequences, such as those polynucleotides in Table 1.

The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the
30 terms "stringent conditions" and "stringent hybridization conditions" mean hybridization occurring only if there is at least 95% and preferably at least 97% identity between the sequences. A specific

example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 or 3 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or 3 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding MurC and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the MurC gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

A coding region of a MurC gene may be isolated by screening using a DNA sequence provided in Table 1 [SEQ ID NO: 1 or 3] to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

There are several methods available and well known to those skilled in the art to obtain full-length DNAs, or extend short DNAs, for example those based on the method of Rapid

Amplification of cDNA ends (RACE) (see, for example, Frohman, *et al.*, *PNAS USA* 85: 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the DNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the selected gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length DNA constructed either by joining the product directly to the existing DNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for diseases, particularly human diseases, as further discussed herein relating to polynucleotide assays.

The polynucleotides of the invention that are oligonucleotides derived from a sequence of Table 1 [SEQ ID NOS:1 or 2 or 3 or 4] may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

For each and every polynucleotide of the invention there is provided a polynucleotide complementary to it. It is preferred that these complementary polynucleotides are fully complementary to each polynucleotide with which they are complementary.

A precursor protein, having a mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

5 In addition to the standard A, G, C, T/U representations for nucleotides, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA nucleotides may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a nucleic acid that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of
10 generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more
15 prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

Vectors, Host Cells, Expression Systems

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the
20 production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression
25 systems. Accordingly, in a further aspect, the present invention relates to expression systems which comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

For recombinant production of the polypeptides of the invention, host cells can be
30 genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods

described in many standard laboratory manuals, such as Davis, *et al.*, *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci *E. coli*, streptomyces, cyanobacteria, *Bacillus subtilis*, and *Staphylococcus aureus*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picornaviruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion

or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic, Prognostic, Serotyping and Mutation Assays

This invention is also related to the use of MurC polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of MurC polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the MurC gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.

Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from a related organism, preferably a different species of the same genus or a different strain of the same species. Point mutations can be identified by hybridizing amplified DNA to labeled MurC polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers *et al.*, *Science*, 230: 1242 (1985). Sequence changes at specific locations also

may be revealed by nuclease protection assays, such as RNase, V1 and S1 protection assay or a chemical cleavage method. See, for example, Cotton *et al.*, *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

In another embodiment, an array of oligonucleotides probes comprising MurC nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype, taxonomic classification or identification. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee *et al.*, *Science*, 274: 610 (1996)).

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1 or 3, or a fragment thereof ;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or 4 or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2 or 4.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a Disease, among others.

This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention, preferable, SEQ ID NO:1 or 3, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease, which results from under-expression, over-expression or altered expression of the polynucleotide. Organisms, particularly infectious organisms, carrying mutations in such polynucleotide may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.

The nucleotide sequences of the present invention are also valuable for organism chromosome identification. The sequence is specifically targeted to, and can hybridize with, a particular location on an organism's chromosome, particularly to a *Staphylococcus aureus*

chromosome. The mapping of relevant sequences to chromosomes according to the present invention may be an important step in correlating those sequences with pathogenic potential and/or an ecological niche of an organism and/or drug resistance of an organism, as well as the essentiality of the gene to the organism. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data may be found on-line in a sequence database. The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through known genetic methods, for example, through linkage analysis (coinheritance of physically adjacent genes) or mating studies, such as by conjugation.

The differences in a polynucleotide and/or polypeptide sequence between organisms possessing a first phenotype and organisms possessing a different, second different phenotype can also be determined. If a mutation is observed in some or all organisms possessing the first phenotype but not in any organisms possessing the second phenotype, then the mutation is likely to be the causative agent of the first phenotype.


Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR. As an example, PCR primers complementary to a polynucleotide encoding MurC polypeptide can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

Table 2

Primers for amplification of MurC polynucleotides

<u>SEQ ID NO</u>	<u>PRIMER SEQUENCE</u>
5	5' -CTTCATTAATGAACGATGC-3'
6	5' -GTTACAAATATTAAAGAAG-3'

The invention also includes primers of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, a metal or a modified nucleotide residue, and at the 3' end of the molecule, Y is hydrogen, a metal or a modified nucleotide residue, R₁ and R₃ are any nucleic acid residue or modified nucleotide residue, m is an integer between 1 and 20 or zero, n is an integer between 1 and 20 or zero, and R₂ is a primer sequence of the invention, particularly a primer sequence selected from Table 2. In the polynucleotide formula above R₂ is oriented so that its 5' end nucleotide residue is at the left, bound to R₁, and its 3' end nucleotide residue is at the right, bound to R₃. Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer being complementary to a region of a polynucleotide of Table 1. In a preferred embodiment m and/or n is an integer between 1 and 10.

The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying MurC DNA and/or RNA isolated from a sample derived from an individual, such as a bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections caused by *Staphylococcus aureus*, comprising determining from a sample derived from an individual, such as a bodily material, an increased level of expression of polynucleotide having a sequence of Table 1 [SEQ ID NO: 1 or 3]. Increased or decreased expression of a MurC polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting, spectrometry and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of MurC polypeptide compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a MurC polypeptide, in a sample derived from a host, such as a bodily material, are well-known to

those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich assays, antibody detection and ELISA assays.

Differential Expression

The polynucleotides and polynucleotides of the invention may be used as reagents for differential screening methods. There are many differential screening and differential display methods known in the art in which the polynucleotides and polypeptides of the invention may be used. For example, the differential display technique is described by Chuang *et al.*, *J. Bacteriol.* 175:2026-2036 (1993). This method identifies those genes which are expressed in an organism by identifying mRNA present using randomly-primed RT-PCR. By comparing pre-infection and post infection profiles, genes up and down regulated during infection can be identified and the RT-PCR product sequenced and matched to ORF "unknowns."

In Vivo Expression Technology (IVET) is described by Camilli *et al.*, *Proc. Nat'l. Acad. Sci. USA.* 91:2634-2638 (1994). IVET identifies genes up-regulated during infection when compared to laboratory cultivation, implying an important role in infection. ORFs identified by this technique are implied to have a significant role in infection establishment and/or maintenance. In this technique random chromosomal fragments of target organism are cloned upstream of a promoter-less recombinase gene in a plasmid vector. This construct is introduced into the target organism which carries an antibiotic resistance gene flanked by resolvase sites. Growth in the presence of the antibiotic removes from the population those fragments cloned into the plasmid vector capable of supporting transcription of the recombinase gene and therefore have caused loss of antibiotic resistance. The resistant pool is introduced into a host and at various times after infection bacteria may be recovered and assessed for the presence of antibiotic resistance. The chromosomal fragment carried by each antibiotic sensitive bacterium should carry a promoter or portion of a gene normally upregulated during infection. Sequencing upstream of the recombinase gene allows identification of the up regulated gene.

RT-PCR may also be used to analyze gene expression patterns. For RT PCR using the polynucleotides of the invention, messenger RNA is isolated from bacterial infected tissue, e.g., 48 hour murine lung infections, and the amount of each mRNA species assessed by reverse transcription of the RNA sample primed with random hexanucleotides followed by PCR with gene specific primer pairs. The determination of the presence and amount of a particular mRNA species by quantification of the resultant PCR product provides information on the bacterial

genes which are transcribed in the infected tissue. Analysis of gene transcription can be carried out at different times of infection to gain a detailed knowledge of gene regulation in bacterial pathogenesis allowing for a clearer understanding of which gene products represent targets for screens for antibacterials. Because of the gene specific nature of the PCR primers employed it should be understood that the bacterial mRNA preparation need not be free of mammalian RNA. This allows the investigator to carry out a simple and quick RNA preparation from infected tissue to obtain bacterial mRNA species which are very short lived in the bacterium (in the order of 2 minute halflives). Optimally the bacterial mRNA is prepared from infected murine lung tissue by mechanical disruption in the presence of TRIzole (GIBCO-BRL) for very short periods of time, subsequent processing according to the manufacturers of TRIzole reagent and DNAase treatment to remove contaminating DNA. Preferably the process is optimized by finding those conditions which give a maximum amount of *Staphylococcus aureus* 16S ribosomal RNA as detected by probing Northernblots with a suitably labeled sequence specific oligonucleotide probe. Typically a 5' dye labeled primer is used in each PCR primer pair in a PCR reaction which is terminated optimally between 8 and 25 cycles. The PCR products are separated on 6% polyacrylamide gels with detection and quantification using GeneScanner (manufactured by ABI).

Gridding and Polynucleotide Subtraction

Methods have been described for obtaining information about gene expression and identity using so called "high density DNA arrays" or grids. See, e.g., M. Chee *et al.*, *Science*, 274:610-614 (1996) and other references cited therein. Such gridding assays have been employed to identify certain novel gene sequences, referred to as Expressed Sequence Tags (EST) (*Adams et al.*, *Science*, 252:1651-1656 (1991)). A variety of techniques have also been described for identifying particular gene sequences on the basis of their gene products. For example, see International Patent Application No. WO91/07087, published May 30, 1991. In addition, methods have been described for the amplification of desired sequences. For example, see International Patent Application No. WO91/17271, published November 14, 1991.

The polynucleotides of the invention may be used as components of polynucleotide arrays, preferably high density arrays or grids. These high density arrays are particularly useful for diagnostic and prognostic purposes. For example, a set of spots each comprising a different

gene, and further comprising a polynucleotide or polynucleotides of the invention, may be used for probing, such as using hybridization or nucleic acid amplification, using a probes obtained or derived from a bodily sample, to determine the presence of a particular polynucleotide sequence or related sequence in an individual. Such a presence may indicate the presence of a pathogen, particularly *Staphylococcus aureus*, and may be useful in diagnosing and/or prognosing disease or a course of disease. A grid comprising a number of variants of the polynucleotide sequence of SEQ ID NO:1 or 3 are preferred. Also preferred is a comprising a number of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2 or 4.

Antibodies

The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively.

In certain preferred embodiments of the invention there are provided antibodies against MurC polypeptides or polynucleotides.

Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

Alternatively, phage display technology may be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-MurC or from naive

libraries (McCafferty, *et al.*, (1990), *Nature* 348, 552-554; Marks, *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson *et al.*, (1991) *Nature* 352: 628).

5 The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or polynucleotides by, for example, affinity chromatography.

Thus, among others, antibodies against MurC-polypeptide or MurC-polynucleotide may be employed to treat infections, particularly bacterial infections.

10 Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

A polypeptide or polynucleotide of the invention, such as an antigenically or immunologically equivalent derivative or a fusion protein of the polypeptide is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin, keyhole limpet haemocyanin or tetanus toxoid. Alternatively, a multiple antigenic polypeptide comprising multiple copies of the polypeptide, or an antigenically or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

20 Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones *et al.* (1986), *Nature* 321, 522-525 or Tempest *et al.*, (1991) *Biotechnology* 9, 266-273.

25 In accordance with an aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally occurring allelic variants of MurC polynucleotides and polypeptides encoded thereby.

30 The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff *et al.*, *Hum Mol Genet* (1992) 1: 363, Manthorpe *et al.*, *Hum. Gene Ther.* (1983) 4: 419), delivery of DNA complexed with specific protein carriers (Wu *et al.*, *J Biol Chem.* (1989) 264: 16985),

coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, (1986) 83: 9551), encapsulation of DNA in various forms of liposomes (Kaneda *et al.*, *Science* (1989) 243: 375), particle bombardment (Tang *et al.*, *Nature* (1992) 356:152, Eisenbraun *et al.*, *DNA Cell Biol* (1993) 12: 791) and *in vivo* infection using cloned retroviral vectors (Seeger *et al.*, *PNAS USA* (1984) 81: 5849).

Antagonists and Agonists - Assays and Molecules

Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, *e.g.*, Coligan *et al.*, *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

Polypeptides and polynucleotides of the present invention are responsible for many biological functions, including many disease states, in particular the Diseases hereinbefore mentioned. It is therefore desirable to devise screening methods to identify compounds which stimulate or which inhibit the function of the polypeptide or polynucleotide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those which stimulate or which inhibit the function of a polypeptide or polynucleotide of the invention, as well as related polypeptides and polynucleotides. In general, agonists or antagonists may be employed for therapeutic and prophylactic purposes for such Diseases as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. Such agonists, antagonists or inhibitors so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of MurC polypeptides and polynucleotides; or may be structural or functional mimetics thereof (see Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991)).

The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or

polynucleotide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring MurC polypeptide and/or polynucleotide activity in the mixture, and comparing the MurC polypeptide and/or polynucleotide activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and MurC polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and and/or functionally related polypeptides (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

The polynucleotides, polypeptides and antibodies that bind to and/or interact with a polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and/or polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of MurC polypeptides or polynucleotides, particularly those compounds that are bacteristatic and/or bactericidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising MurC polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a MurC agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the MurC polypeptide is reflected in decreased binding of the labeled ligand or decreased production of

product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of MurC polypeptide are most likely to be good antagonists. Molecules that bind well and, as the case may be, increase the rate of product production from substrate, increase signal transduction, or increase chemical channel activity are agonists. Detection of the rate or level of, as the case may be, production of product from substrate, signal transduction, or chemical channel activity may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric, labeled substrate converted into product, a reporter gene that is responsive to changes in MurC polynucleotide or polypeptide activity, and binding assays known in the art.

Polypeptides of the invention may be used to identify membrane bound or soluble receptors, if any, for such polypeptide, through standard receptor binding techniques known in the art. These techniques include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance, ^{125}I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (*e.g.*, cells, cell membranes, cell supernatants, tissue extracts, bodily materials). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide which compete with the binding of the polypeptide to its receptor(s), if any. Standard methods for conducting such assays are well understood in the art.

The fluorescence polarization value for a fluorescently-tagged molecule depends on the rotational correlation time or tumbling rate. Protein complexes, such as formed by MurC polypeptide associating with another MurC polypeptide or other polypeptide, labeled to comprise a fluorescently-labeled molecule will have higher polarization values than a fluorescently labeled monomeric protein. It is preferred that this method be used to characterize small molecules that disrupt polypeptide complexes.

Fluorescence energy transfer may also be used characterize small molecules that interfere with the formation of MurC polypeptide dimers, trimers, tetramers or higher order structures, or structures formed by MurC polypeptide bound to another polypeptide. MurC polypeptide can be labeled with both a donor and acceptor fluorophore. Upon mixing of the two labeled species and excitation of the donor fluorophore, fluorescence energy transfer can

be detected by observing fluorescence of the acceptor. Compounds that block dimerization will inhibit fluorescence energy transfer.

Surface plasmon resonance can be used to monitor the effect of small molecules on MurC polypeptide self-association as well as an association of MurC polypeptide and another polypeptide or small molecule. MurC polypeptide can be coupled to a sensor chip at low site density such that covalently bound molecules will be monomeric. Solution protein can then be passed over the MurC polypeptide-coated surface and specific binding can be detected in real-time by monitoring the change in resonance angle caused by a change in local refractive index. This technique can be used to characterize the effect of small molecules on kinetic rates and equilibrium binding constants for MurC polypeptide self-association as well as an association of MurC polypeptide and another polypeptide or small molecule.

A scintillation proximity assay may be used to characterize the interaction between an association of MurC polypeptide with another MurC polypeptide or a different polypeptide. MurC polypeptide can be coupled to a scintillation-filled bead. Addition of radio-labeled MurC polypeptide results in binding where the radioactive source molecule is in close proximity to the scintillation fluid. Thus, signal is emitted upon MurC polypeptide binding and compounds that prevent MurC polypeptide self-association or an association of MurC polypeptide and another polypeptide or small molecule will diminish signal.

ICS biosensors have been described by AMBRI (Australian Membrane Biotechnology Research Institute). They couple the self-association of macromolecules to the closing of gramicidin-facilitated ion channels in suspended membrane bilayers and hence to a measurable change in the admittance (similar to impedance) of the biosensor. This approach is linear over six decades of admittance change and is ideally suited for large scale, high through-put screening of small molecule combinatorial libraries.

In other embodiments of the invention there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity or expression of a polypeptide and/or polynucleotide of the invention comprising: contacting a polypeptide and/or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other interaction between the compound and the polypeptide and/or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction preferably being associated with a second component capable of providing a detectable signal in response to the

binding or interaction of the polypeptide and/or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity or expression of the polypeptide and/or polynucleotide by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide and/or polynucleotide.

Another example of an assay for MurC agonists is a competitive assay that combines MurC and a potential agonist with MurC-binding molecules, recombinant MurC binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. MurC can be labeled, such as by radioactivity or a colorimetric compound, such that the number of MurC molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing MurC-induced activities, thereby preventing the action or expression of MurC polypeptides and/or polynucleotides by excluding MurC polypeptides and/or polynucleotides from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of MurC.

Other examples of potential polypeptide antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or small molecules which bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Certain of the polypeptides of the invention are biomimetics, functional mimetics of the natural MurC polypeptide. These functional mimetics may be used for, among other things, antagonizing the activity of MurC polypeptide or as a antigen or immunogen in a manner described elsewhere herein. Functional mimetics of the polypeptides of the invention include but are not limited to truncated polypeptides. For example, preferred functional mimetics include, a polypeptide comprising the polypeptide sequence set forth in SEQ ID NO:2 lacking 20, 30, 40, 50, 60, 70 or 80 amino- or carboxy-terminal amino acid residues, including fusion proteins comprising one or more of these truncated sequences. Polynucleotides encoding each of these functional mimetics may be used as expression cassettes to express each mimetic polypeptide. It is preferred that these cassettes comprise 5' and 3' restriction sites to allow for a convenient means to ligate the cassettes together when desired. It is further preferred that these cassettes comprise gene expression signals known in the art or described elsewhere herein.

Thus, in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for a polypeptide and/or polynucleotide of the present invention; or compounds which decrease or enhance the production of such polypeptides and/or polynucleotides, which comprises:

- (a) a polypeptide and/or a polynucleotide of the present invention;
 - (b) a recombinant cell expressing a polypeptide and/or polynucleotide of the present invention;
 - (c) a cell membrane expressing a polypeptide and/or polynucleotide of the present invention; or
 - (d) antibody to a polypeptide and/or polynucleotide of the present invention;
- which polypeptide is preferably that of SEQ ID NO:2, and which polynucleotide is preferably that of SEQ ID NO:1.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

It will be readily appreciated by the skilled artisan that a polypeptide and/or polynucleotide of the present invention may also be used in a method for the structure-based design of an agonist, antagonist or inhibitor of the polypeptide and/or polynucleotide, by:

- (a) determining in the first instance the three-dimensional structure of the polypeptide and/or polynucleotide, or complexes thereof;
- (b) deducing the three-dimensional structure for the likely reactive site(s), binding site(s) or motif(s) of an agonist, antagonist or inhibitor;

(c) synthesizing candidate compounds that are predicted to bind to or react with the deduced binding site(s), reactive site(s), and/or motif(s); and

(d) testing whether the candidate compounds are indeed agonists, antagonists or inhibitors.

It will be further appreciated that this will normally be an iterative process, and this iterative

5 process may be performed using automated and computer-controlled steps.

In a further aspect, the present invention provides methods of treating abnormal conditions such as, for instance, a Disease, related to either an excess of, an under-expression of, an elevated activity of, or a decreased activity of MurC polypeptide and/or polynucleotide.

10 If the expression and/or activity of the polypeptide and/or polynucleotide is in excess, several approaches are available. One approach comprises administering to an individual in need thereof an inhibitor compound (antagonist) as herein described, optionally in combination with a pharmaceutically acceptable carrier, in an amount effective to inhibit the function and/or expression of the polypeptide and/or polynucleotide, such as, for example, by blocking the binding of ligands, substrates, receptors, enzymes, etc., or by inhibiting a second signal, and thereby alleviating the

15 abnormal condition. In another approach, soluble forms of the polypeptides still capable of binding the ligand, substrate, enzymes, receptors, etc. in competition with endogenous polypeptide and/or polynucleotide may be administered. Typical examples of such competitors include fragments of the MurC polypeptide and/or polypeptide.

In a further aspect, the present invention relates to genetically engineered soluble fusion

20 proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage

25 sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

30 In still another approach, expression of the gene encoding endogenous MurC polypeptide can be inhibited using expression blocking techniques. This blocking may be targeted against

any step in gene expression, but is preferably targeted against transcription and/or translation. An examples of a known technique of this sort involve the use of antisense sequences, either internally generated or separately administered (see, for example, O'Connor, *J Neurochem* (1991) 56:560 in *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices with the gene can be supplied (see, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360). These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*.

Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to interfere with the initial physical interaction between a pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial MurC proteins that mediate tissue damage and/or; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

In accordance with yet another aspect of the invention, there are provided MurC agonists and antagonists, preferably bacteristatic or bactericidal agonists and antagonists.

The antagonists and agonists of the invention may be employed, for instance, to prevent, inhibit and/or treat diseases.

Helicobacter pylori (herein "*H. pylori*") bacteria infect the stomachs of over one-third of the world's population causing stomach cancer, ulcers, and gastritis (International Agency for Research on Cancer (1994) *Schistosomes, Liver Flukes and Helicobacter Pylori* (International

Agency for Research on Cancer, Lyon, France, <http://www.uicc.ch/ecp/ecp2904.htm>). Moreover, the International Agency for Research on Cancer recently recognized a cause-and-effect relationship between *H. pylori* and gastric adenocarcinoma, classifying the bacterium as a Group I (definite) carcinogen. Preferred antimicrobial compounds of the invention (agonists and antagonists of MurC polypeptides and/or polynucleotides) found using screens provided by the invention, or known in the art, particularly narrow-spectrum antibiotics, should be useful in the treatment of *H. pylori* infection. Such treatment should decrease the advent of *H. pylori*-induced cancers, such as gastrointestinal carcinoma. Such treatment should also prevent, inhibit and/or cure gastric ulcers and gastritis.

Vaccines

There are provided by the invention, products, compositions and methods for assessing MurC expression, treating disease, assaying genetic variation, and administering a MurC polypeptide and/or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Staphylococcus aureus* bacteria.

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with MurC polynucleotide and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Staphylococcus aureus* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct expression of MurC polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing MurC polynucleotide and/or polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease, whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

5 A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a MurC polynucleotide and/or polypeptide encoded therefrom, wherein the composition comprises a recombinant MurC polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said MurC polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity and/or cellular immunity, such as cellular immunity arising from CTL or CD4+ T cells.

10 A MurC polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of stabilizing the first protein and producing a fused or modified protein which will have antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus*
15 *influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and facilitates production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

20 Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

25 Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Staphylococcus aureus*. Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of
30 monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or

therapeutic treatments of bacterial infection, particularly *Staphylococcus aureus* infection, in mammals, particularly humans.

5 A polypeptide of the invention may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused, for example, by mechanical, chemical, thermal or radiation damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, throat, mammary glands, urethra or vagina.

10 The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile
15 injection solutions which may contain anti-oxidants, buffers, bacteristatic compounds and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition
20 requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

25 While the invention has been described with reference to certain MurC polypeptides and polynucleotides, it is to be understood that this covers fragments of the naturally occurring polypeptides and polynucleotides, and similar polypeptides and polynucleotides with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant polypeptides or polynucleotides.

30 **Compositions, kits and administration**

051033287 "062398
062398
In a further aspect of the invention there are provided compositions comprising a MurC polynucleotide and/or a MurC polypeptide for administration to a cell or to a multicellular organism.

5 The invention also relates to compositions comprising a polynucleotide and/or a polypeptides discussed herein or their agonists or antagonists. The polypeptides and polynucleotides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to an individual. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide and/or polynucleotide of the invention and a
10 pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

15 Polypeptides, polynucleotides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

20 In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate
25 conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

30 In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide and/or polynucleotide, such as the

soluble form of a polypeptide and/or polynucleotide of the present invention, agonist or antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the
5 aforementioned compositions of the invention. Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The composition will be adapted to the route of administration, for instance by a systemic
10 or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be
15 formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, and the like.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg.
20 The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e.,
25 devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a systemic
30 effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition, the composition

could also be used to broaden perioperative cover for any surgical technique to prevent bacterial wound infections, especially *Staphylococcus aureus* wound infections.

Many orthopedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia.

- 5 Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

- 10 In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

- 15 A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

20 **Sequence Databases, Sequences in a Tangible Medium, and Algorithms**

- Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as GCG.

- 25 The polynucleotide and polypeptide sequences of the invention are particularly useful as components in databases useful for search analyses as well as in sequence analysis algorithms. As used in this section entitled "Sequence Databases, Sequences in a Tangible Medium, and Algorithms," and in claims related to this section, the terms "polynucleotide of the invention" and "polynucleotide sequence of the invention" mean any detectable chemical or physical

characteristic of a polynucleotide of the invention that is or may be reduced to or stored in a tangible medium, preferably a computer readable form. For example, chromatographic scan data or peak data, photographic data or scan data therefrom, called bases, and mass spectrographic data. As used in this section entitled Databases and Algorithms and in claims related thereto, the

5 terms "polypeptide of the invention" and "polypeptide sequence of the invention" mean any detectable chemical or physical characteristic of a polypeptide of the invention that is or may be reduced to or stored in a tangible medium, preferably a computer readable form. For example, chromatographic scan data or peak data, photographic data or scan data therefrom, and mass spectrographic data.

10 The invention provides a computer readable medium having stored thereon polypeptide sequences of the invention and/or polynucleotide sequences of the invention. For example, a computer readable medium is provided comprising and having stored thereon a member selected from the group consisting of: a polynucleotide comprising the sequence of a polynucleotide of the invention; a polypeptide comprising the sequence of a polypeptide sequence of the invention;

15 a set of polynucleotide sequences wherein at least one of the sequences comprises the sequence of a polynucleotide sequence of the invention; a set of polypeptide sequences wherein at least one of the sequences comprises the sequence of a polypeptide sequence of the invention; a data set representing a polynucleotide sequence comprising the sequence of polynucleotide sequence of the invention; a data set representing a polynucleotide sequence encoding a polypeptide sequence

20 comprising the sequence of a polypeptide sequence of the invention; a polynucleotide comprising the sequence of a polynucleotide sequence of the invention; a polypeptide comprising the sequence of a polypeptide sequence of the invention; a set of polynucleotide sequences wherein at least one of the sequences comprises the sequence of a polynucleotide sequence of the invention; a set of polypeptide sequences wherein at least one of said sequences comprises the

25 sequence of a polypeptide sequence of the invention; a data set representing a polynucleotide sequence comprising the sequence of a polynucleotide sequence of the invention; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of a polypeptide sequence of the invention. The computer readable medium can be any composition of matter used to store information or data, including, for example, commercially

30 available floppy disks, tapes, chips, hard drives, compact disks, and video disks.

Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded genetic sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, RNA structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination, nucleic acid base calling, nucleic acid base trimming, and sequencing chromatogram peak analysis.

A computer based method is provided for performing homology identification. This method comprises the steps of providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

A computer based method is still further provided for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and screening for at least one overlapping region between said first polynucleotide sequence and at least one second polynucleotide or polypeptide sequence.

A computer based method is still further provided for polynucleotide assembly, said method comprising the steps of: providing a first polypeptide sequence comprising a polypeptide of the invention in a computer readable medium; and screening for at least one overlapping region between said first polypeptide sequence and at least one second polynucleotide or polypeptide sequence.

In another preferred embodiment of the invention there is provided a computer readable medium having stored thereon a member selected from the group consisting of: a polynucleotide comprising the sequence of SEQ ID NO. 1 or 3; a polypeptide comprising the sequence of SEQ ID NO. 2 or 4; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 1 or 3; a set of polypeptide sequences wherein at least

one of said sequences comprises the sequence of SEQ ID NO. 2 or 4; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2 or 4; a polynucleotide comprising the sequence of SEQ ID NO. 1 or 3; a polypeptide comprising the sequence of SEQ ID NO. 2 or 4; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 1 or 3; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 2 or 4; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2 or 4. A further preferred embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of providing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3 in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

15 A still further preferred embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of: providing a polypeptide sequence comprising the sequence of SEQ ID NO. 2 or 4 in a computer readable medium; and comparing said polypeptide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

20 A further embodiment of the invention provides a computer based method for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3 in a computer readable medium; and screening for at least one overlapping region between said first polynucleotide sequence and a second polynucleotide sequence.

25 A further embodiment of the invention provides a computer based method for performing homology identification, said method comprising the steps of: providing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3 in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

30 All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if

each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

5

GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

10 "Antibody(ies)" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Antigenically equivalent derivative(s)" as used herein encompasses a polypeptide, polynucleotide, or the equivalent of either which will be specifically recognized by certain antibodies which, when raised to the protein, polypeptide or polynucleotide according to the
15 invention, interferes with the immediate physical interaction between pathogen and mammalian host.

"Bispecific antibody(ies)" means an antibody comprising at least two antigen binding domains, each domain directed against a different epitope.

20 "Bodily material(s)" means any material derived from an individual or from an organism infecting, infesting or inhabiting an individual, including but not limited to, cells, tissues and waste, such as, bone, blood, serum, cerebrospinal fluid, semen, saliva, muscle, cartilage, organ tissue, skin, urine, stool or autopsy materials..

"Disease(s)" means any disease caused by or related to infection by a bacteria, including , for example, disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac
25 (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic absces, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric absces, toxic shock syndrome), skin (e.g., impetigo,
30 folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis).

"Fusion protein(s)" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0464 discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties [see, e.g., EP-A 0232262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified.

"Host cell(s)" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). The well known Smith Waterman algorithm may also be used to determine identity.

Parameters for polypeptide sequence comparison include the following:

1) Algorithm: Needleman and Wunsch, J. Mol Biol. 48: 443-453 (1970)

Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992)

Gap Penalty: 12

5 Gap Length Penalty: 4

A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

10 1) Algorithm: Needleman and Wunsch, J. Mol Biol. 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

15 Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the default parameters for nucleic acid comparisons.

A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

- (1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO: 1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \bullet y),$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \bullet y),$$

wherein n_n is the number of nucleic acid alterations, x_n is the total number of nucleic acids in SEQ ID NO:1, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

09103237-062398
055399-49230760

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50,60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO: 2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

15
$$n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of

amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

5
$$n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

"Immunologically equivalent derivative(s)" as used herein encompasses a polypeptide, polynucleotide, or the equivalent of either which when used in a suitable formulation to raise antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

15 "Immunospecific" means that characteristic of an antibody whereby it possesses substantially greater affinity for the polypeptides of the invention or the polynucleotides of the invention than its affinity for other related polypeptides or polynucleotides respectively, particularly those polypeptides and polynucleotides in the prior art.

"Individual(s)" means a multicellular eukaryote, including, but not limited to a metazoan, a mammal, an ovid, a bovid, a simian, a primate, and a human.

20 "Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

25 "Organism(s)" means a (i) prokaryote, including but not limited to, a member of the genus *Streptococcus*, *Staphylococcus*, *Bordetella*, *Corynebacterium*, *Mycobacterium*, *Neisseria*, *Haemophilus*, *Actinomycetes*, *Streptomyces*, *Nocardia*, *Enterobacter*, *Yersinia*, *Fancisella*,

Pasturella, *Moraxella*, *Acinetobacter*, *Erysipelothrix*, *Branhamella*, *Actinobacillus*, *Streptobacillus*, *Listeria*, *Calymmatobacterium*, *Brucella*, *Bacillus*, *Clostridium*, *Treponema*, *Escherichia*, *Salmonella*, *Klebsiella*, *Vibrio*, *Proteus*, *Erwinia*, *Borrelia*, *Leptospira*, *Spirillum*, *Campylobacter*, *Shigella*, *Legionella*, *Pseudomonas*, *Aeromonas*, *Rickettsia*, *Chlamydia*, *Borrelia* and *Mycoplasma*,

- 5 and further including, but not limited to, a member of the species or group, Group A *Streptococcus*, Group B *Streptococcus*, Group C *Streptococcus*, Group D *Streptococcus*, Group G *Streptococcus*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus faecalis*, *Streptococcus faecium*, *Streptococcus durans*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Corynebacterium diphtheriae*,
10 *Gardnerella vaginalis*, *Mycobacterium tuberculosis*, *Mycobacterium bovis*, *Mycobacterium ulcerans*, *Mycobacterium leprae*, *Actinomyces israelii*, *Listeria monocytogenes*, *Bordetella pertussis*, *Bordetella parapertussis*, *Bordetella bronchiseptica*, *Escherichia coli*, *Shigella dysenteriae*, *Haemophilus influenzae*, *Haemophilus aegyptius*, *Haemophilus parainfluenzae*, *Haemophilus ducreyi*, *Bordetella*, *Salmonella typhi*, *Citrobacter freundii*, *Proteus mirabilis*, *Proteus vulgaris*,
15 *Yersinia pestis*, *Klebsiella pneumoniae*, *Serratia marcescens*, *Serratia liquefaciens*, *Vibrio cholera*, *Shigella dysenteriae*, *Shigella flexneri*, *Pseudomonas aeruginosa*, *Francisella tularensis*, *Brucella abortus*, *Bacillus anthracis*, *Bacillus cereus*, *Clostridium perfringens*, *Clostridium tetani*, *Clostridium botulinum*, *Treponema pallidum*, *Rickettsia rickettsii* and *Chlamydia trachomatis*, (ii) an archaeon, including but not limited to *Archaeobacter*, and (iii) a unicellular or filamentous eukaryote,
20 including but not limited to, a protozoan, a fungus, a member of the genus *Saccharomyces*, *Kluyveromyces*, or *Candida*, and a member of the species *Saccharomyces cerevisiae*, *Kluyveromyces lactis*, or *Candida albicans*.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include,
25 without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used
30 herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions

may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing,

phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

"Recombinant expression system(s)" refers to expression systems or portions thereof or polynucleotides of the invention introduced or transformed into a host cell or host cell lysate for the production of the polynucleotides and polypeptides of the invention.

"Subtraction set" is one or more, but preferably less than 100, polynucleotides comprising at least one polynucleotide of the invention

"Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusion proteins and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. The present invention also includes include variants of each of the polypeptides of the invention, that is polypeptides that vary from the referents by conservative

amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

10 **EXAMPLES**

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1 Strain selection, Library Production and Sequencing

15 The polynucleotide having a DNA sequence given in Table 1 [SEQ ID NO:1 or 3] was obtained from a library of clones of chromosomal DNA of *Staphylococcus aureus* in *E. coli*. The sequencing data from two or more clones containing overlapping *Staphylococcus aureus* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

20 Methods 1 and 2 below.

Total cellular DNA is isolated from *Staphylococcus aureus* WCUH 29 according to standard procedures and size-fractionated by either of two methods.

Method 1

25 Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

30 **Method 2**

Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, BshI235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Example 2 MurC Characterization

Determination of expression, during infection, of a gene from *Staphylococcus aureus*

Necrotic fatty tissue from a four day groin infection of *Staphylococcus aureus* WCUH29 in the mouse is efficiently disrupted and processed in the presence of chaotropic agents and RNAase inhibitor to provide a mixture of animal and bacterial RNA. The optimal conditions for disruption and processing to give stable preparations and high yields of bacterial RNA are followed by the use of hybridisation to a radiolabelled oligonucleotide specific to *Staphylococcus aureus* 16S RNA on Northern blots. The RNAase free, DNAase free, DNA and protein free preparations of RNA obtained are suitable for Reverse Transcription PCR (RT-PCR) using unique primer pairs designed from the sequence of each gene of *Staphylococcus aureus* WCUH29.

a) Isolation of tissue infected with *Staphylococcus aureus* WCUH29 from a mouse animal model of infection

10 ml. volumes of sterile nutrient broth (No.2 Oxoid) are seeded with isolated, individual colonies of *Staphylococcus aureus* WCUH29 from an agar culture plate. The cultures are incubated aerobically (static culture) at 37°C for 16-20 hours. 4 week old mice (female, 18g-22g, strain MF1) are each infected by subcutaneous injection of 0.5ml. of this broth culture of *Staphylococcus aureus* WCUH29 (diluted in broth to approximately 10⁸ cfu/ml.) into the anterior, right lower quadrant (groin area). Mice should be monitored regularly during the first 24 hours after infection, then daily until termination of study. Animals with signs of systemic infection, i.e. lethargy, ruffled appearance, isolation from group, should be monitored closely and if signs progress to moribundancy, the animal should be culled immediately.

Visible external signs of lesion development will be seen 24-48h after infection. Examination of the abdomen of the animal will show the raised outline of the abscess beneath the skin. The localised lesion should remain in the right lower quadrant, but may occasionally spread

to the left lower quadrant, and superiorly to the thorax. On occasions, the abscess may rupture through the overlying skin layers. In such cases the affected animal should be culled immediately and the tissues sampled if possible. Failure to cull the animal may result in the necrotic skin tissue overlying the abscess sloughing off, exposing the abdominal muscle wall.

5 Approximately 96 hours after infection, animals are killed using carbon dioxide asphyxiation. To minimise delay between death and tissue processing /storage, mice should be killed individually rather than in groups. The dead animal is placed onto its back and the fur swabbed liberally with 70% alcohol. An initial incision using scissors is made through the skin of the abdominal left lower quadrant, travelling superiorly up to, then across the thorax. The
10 incision is completed by cutting inferiorly to the abdominal lower right quadrant. Care should be taken not to penetrate the abdominal wall. Holding the skin flap with forceps, the skin is gently pulled away from the abdomen. The exposed abscess, which covers the peritoneal wall but generally does not penetrate the muscle sheet completely, is excised, taking care not to puncture the viscera

15 The abscess/muscle sheet and other infected tissue may require cutting in sections, prior to flash-freezing in liquid nitrogen, thereby allowing easier storage in plastic collecting vials.

b) Isolation of *Staphylococcus aureus* WCUH29 RNA from infected tissue samples

20 4-6 infected tissue samples (each approx 0.5-0.7g) in 2ml screw-cap tubes are removed from -80°C storage into a dry ice ethanol bath. In a microbiological safety cabinet the samples are disrupted individually whilst the remaining samples are kept cold in the dry ice ethanol bath. To disrupt the bacteria within the tissue sample 1ml of TRIzol Reagent (Gibco BRL, Life Technologies) is added followed by enough 0.1mm zirconia/silica beads to almost fill the tube, the lid is replaced taking care not to get any beads into the screw thread so as to ensure a good
25 seal and eliminate aerosol generation. The sample is then homogenised in a Mini-BeadBeater Type BX-4 (Biospec Products). Necrotic fatty tissue is strain treated for 100 seconds at 5000 rpm in order to achieve bacterial lysis. *In vivo* grown bacteria require longer treatment than *in vitro* grown *Staphylococcus aureus* (which are for example disrupted by a 30 second bead-beat).

30 After bead-beating the tubes are chilled on ice before opening in a fume-hood as heat generated during disruption may degrade the TRIzol and release cyanide.

200 microlitres of chloroform is then added and the tubes shaken by hand for 15 seconds to ensure complete mixing. After 2-3 minutes at room temperature the tubes are spun down at 12,000 x g, 4°C for 15 minutes and RNA extraction is then continued according to the method given by the manufacturers of TRIzol Reagent. Specifically, the aqueous phase, approximately 5 0.6 ml, is transferred to a sterile eppendorf tube and 0.5 ml of isopropanol is added. After 10 minutes at room temperature the samples are spun at 12,000 x g, 4°C for 10 minutes. The supernatant is removed and discarded then the RNA pellet is washed with 1 ml 75% ethanol. A brief vortex is used to mix the sample before centrifuging at 7,500 x g, 4°C for 5 minutes. The ethanol is removed and the RNA pellet dried under vacuum for no more than 5 minutes. Samples 10 are then resuspended by repeated pipetting in 100 microlitres of DEPC treated water, followed by 5-10 minutes at 55°C. Finally, after at least 1 minute on ice, 200 units of Rnasin (Promega) is added.

RNA preparations are stored at -80°C for up to one month. For longer term storage the RNA precipitate can be stored at the wash stage of the protocol in 75% ethanol for at least one 15 year at -20°C.

Quality of the RNA isolated is assessed by running samples on 1% agarose gels. 1 x TBE gels stained with ethidium bromide are used to visualise total RNA yields. To demonstrate the isolation of bacterial RNA from the infected tissue 1 x MOPS, 2.2M formaldehyde gels are run and vacuum blotted to Hybond-N (Amersham). The blot is then hybridised with a 32 P labelled 20 oligonucleotide probe specific to 16s rRNA of *Staphylococcus aureus* (K.Greisen, M. Loeffelholz, A. Purohit and D. Leong. J.Clin. (1994) Microbiol. 32 335-351). The size of the hybridising band is compared to that of control RNA isolated from *in vitro* grown *Staphylococcus aureus* WCUH29 in the Northern blot. Correct sized bacterial 16S rRNA bands can be detected in total RNA samples which show extensive degradation of the mammalian RNA when 25 visualised on TBE gels.

c) The removal of DNA from *Staphylococcus aureus* WCUH29-derived RNA

DNA is removed from 73 microlitre samples of RNA by a 15 minute treatment on ice with 3 units of DNAaseI, amplification grade (Gibco BRL, Life Technologies) in the buffer 30 supplied with the addition of 200 units of Rnasin (Promega) in a final volume of 90 microlitres.

The DNAase is inactivated and removed by treatment with TRIzol LS Reagent (Gibco BRL, Life Technologies) according to the manufacturers protocol. DNAase treated RNA is resuspended in 73 microlitres of DEPC treated water with the addition of Rnasin.

5 d) The preparation of cDNA from RNA samples derived from infected tissue

10 microlitre samples of DNAase treated RNA are reverse transcribed using a SuperScript Preamplification System for First Strand cDNA Synthesis kit (Gibco BRL, Life Technologies) according to the manufacturers instructions. 1 nanogram of random hexamers is used to prime each reaction. Controls without the addition of SuperScriptII reverse transcriptase are also run. Both +/-RT samples are treated with RNaseH before proceeding to the PCR reaction

e) The use of PCR to determine the presence of a bacterial cDNA species

15 PCR reactions are set up on ice in 0.2ml tubes by adding the following components: 45 microlitres PCR SUPERMIX (Gibco BRL, Life Technologies); 1 microlitre 50mM MgCl₂ , to adjust final concentration to 2.5mM; 1 microlitre PCR primers (optimally 18-25 basepairs in length and designed to possess similar annealing temperatures), each primer at 10 mM initial concentration; and 2 microlitres cDNA.

20 PCR reactions are run on a Perkin Elmer GeneAmp PCR System 9600 as follows: 5 minutes at 95°C, then 50 cycles of 30 seconds each at 94°C, 42°C and 72°C followed by 3 minutes at 72°C and then a hold temperature of 4°C. (the number of cycles is optimally 30-50 to determine the appearance or lack of a PCR product and optimally 8-30 cycles if an estimation of the starting quantity of cDNA from the RT reaction is to be made); 10 microlitre aliquots are then run out on 1% 1 x TBE gels stained with ethidium bromide with PCR product, if present, sizes estimated by comparison to a 100 bp DNA Ladder (Gibco BRL, Life Technologies).

25 Alternatively if the PCR products are conveniently labelled by the use of a labelled PCR primer (e.g. labelled at the 5'end with a dye) a suitable aliquot of the PCR product is run out on a polyacrylamide sequencing gel and its presence and quantity detected using a suitable gel scanning system (e.g. ABI Prism™ 377 Sequencer using GeneScan™ software as supplied by Perkin Elmer).

RT/PCR controls may include +/- reverse transcriptase reactions, 16S rRNA primers or DNA specific primer pairs designed to produce PCR products from non-transcribed *Staphylococcus aureus* WCUH29 genomic sequences.

To test the efficiency of the primer pairs they are used in DNA PCR with *Staphylococcus aureus* WCUH29 total DNA. PCR reactions are set up and run as described above using approx. 1 microgram of DNA in place of the cDNA and 35 cycles of PCR.

Primer pairs which fail to give the predicted sized product in either DNA PCR or RT/PCR are PCR failures and as such are uninformative. Of those which give the correct size product with DNA PCR two classes are distinguished in RT/PCR: 1. Genes which are not transcribed in vivo reproducibly fail to give a product in RT/PCR; and 2. Genes which are transcribed in vivo reproducibly give the correct size product in RT/PCR and show a stronger signal in the +RT samples than the signal (if at all present) in -RT controls.

The polynucleotide sequence of the invention, SEQ ID NO:1, was identified in the above test as transcribed in vivo in the mouse 7 day pyelonephritis model of infection. SEQ ID NO:2 was deduced from the polynucleotide sequence given as SEQ ID NO:1. The pair of PCR primers used to identify the gene are given as SEQ ID NOS:5 and 6.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Wallis, Nicola G.
Burnham, Martin K. R.

(ii) TITLE OF INVENTION: murC

(iii) NUMBER OF SEQUENCES: 6

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Dechert, Price & Rhoads
(B) STREET: 4000 Bell Atlantic Tower, 1717 Arch Stre
(C) CITY: Philadelphia
(D) STATE: PA
(E) COUNTRY: USA
(F) ZIP: 19103-2793

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: Windows 95
(D) SOFTWARE: FastSEQ for Windows Version 2.0b

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 60/052,720
(B) FILING DATE: 03-JUL-1997

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Falk, Stephen T

- (B) REGISTRATION NUMBER: 36,795
(C) REFERENCE/DOCKET NUMBER: GM10025

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 215-994-2488
(B) TELEFAX: 215-994-2222
(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1351 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGAGTAAGG AGTTTTATAT AATGACACAC TATCATTTTG TCGGAATTAA AGGTTCTGGC 60
ATGAGTTCAT TAGCACAAAT CATGCATGAT TTAGGACATG AAGTTCAAGG ATCGGATATT 120
GAGAACTACG TATTACAGA AGTTGCTCTT AGAAATAAGG GGATAAAAAT ATTACCATTT 180
GGTGCTAATA ACATAAAAGA AGATATGGTA GTTATACAAG GTAATGCATT CGCGAGTAGC 240
CATGAAGAAA TAGTACGTGC ACATCAATTG AAATTAGATG TTGTAAGTTA TAATGATTTT 300
TTAGGACAGA TTATTGATCA ATATACTTCA GTAGCTGTAA CTGGTGCACA TGGTAAACT 360
TCTACAACAG GTTTATTATC ACATGTTATG AATGGTGATA AAAAGACTTC ATTTTTAATT 420
GGTGATGGCA CAGGTATGGG ATTGCCTGAA AGTGATTATT TCGCTTTTGA GGCATGTGAA 480
TATAGACGTC ACTTTTTAAG TTATAAACCT GATTACGCAA TTATGACAAA TATTGATTTT 540
GATCATCCTG ATTATTTCAA AGATATTAAT GATGTTTTTG ATGCATTCCA AGAAATGGCA 600
CATAATGTTA AAAAAGGTAT TATTGCTTGG GGTGATGATG AACATCTACG TAAAATTGAA 660
GCAGATGTTT CAATTTATTA CTATGGATTT AAAGATTCCG ATGACATTTA TGCTCAAAAT 720
ATTCAAATTA CGGATAAAGG TACTGCTTTT GATGTGTATG TGGATGGTGA GTTTTATGAT 780
CACTTCCTGT CTCCACAATA TGGTGACCAT ACAGTTTAA ATGCATTAGC TGTAATTGCG 840
ATTAGTTATT TAGAGAAGCT AGATGTTACA AATATTAAAG AAGCATTAGA AACGTTTGGT 900
GGTGTTAAAC GTCGTTTCAA TGAACTACA ATTGCAAATC AAGTTATTGT AGATGATTAT 960
GCACACCATC CAAGAGAAAT TAGTGCTACA ATTGACACAG CACGAAAGAA ATATCCACAT 1020
AAAGAAGTTG TTGCAGTATT TCAACCACAC ACTTTCTCTA GAACACAAGC ATTTTAAAT 1080
GAATTTGCAG AAAGTTTATG TAAAGCAGAT CGTGTATTCT TATGTGAAAT TTTTGGCTCA 1140
ATTAGAGAAA ATTCTGGCGC ATTAACGATA CAAGATTTAA TTGATAAAAT TGGAGGTGCA 1200
TCGTTCATTA ATGAAGATCT TATTAATGTA TTAGAACAAT TTGATAATGC TGTTGTTTTA 1260

TTTATGGGTG CAGGTGATAT TCAAAAATTA CAAAATGCAT ATTTAGATAA ATTAGGCATG 1320
AAAAATGCGT TTTAATATGT TTATAATAGA G 1351

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 437 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Thr	His	Tyr	His	Phe	Val	Gly	Ile	Lys	Gly	Ser	Gly	Met	Ser	Ser
1				5					10					15	
Leu	Ala	Gln	Ile	Met	His	Asp	Leu	Gly	His	Glu	Val	Gln	Gly	Ser	Asp
			20					25					30		
Ile	Glu	Asn	Tyr	Val	Phe	Thr	Glu	Val	Ala	Leu	Arg	Asn	Lys	Gly	Ile
		35					40					45			
Lys	Ile	Leu	Pro	Phe	Gly	Ala	Asn	Asn	Ile	Lys	Glu	Asp	Met	Val	Val
	50					55					60				
Ile	Gln	Gly	Asn	Ala	Phe	Ala	Ser	Ser	His	Glu	Glu	Ile	Val	Arg	Ala
65					70					75					80
His	Gln	Leu	Lys	Leu	Asp	Val	Val	Ser	Tyr	Asn	Asp	Phe	Leu	Gly	Gln
				85					90					95	
Ile	Ile	Asp	Gln	Tyr	Thr	Ser	Val	Ala	Val	Thr	Gly	Ala	His	Gly	Lys
			100					105					110		
Thr	Ser	Thr	Thr	Gly	Leu	Leu	Ser	His	Val	Met	Asn	Gly	Asp	Lys	Lys
		115					120					125			
Thr	Ser	Phe	Leu	Ile	Gly	Asp	Gly	Thr	Gly	Met	Gly	Leu	Pro	Glu	Ser
	130					135					140				
Asp	Tyr	Phe	Ala	Phe	Glu	Ala	Cys	Glu	Tyr	Arg	Arg	His	Phe	Leu	Ser
145					150					155					160
Tyr	Lys	Pro	Asp	Tyr	Ala	Ile	Met	Thr	Asn	Ile	Asp	Phe	Asp	His	Pro
				165					170					175	
Asp	Tyr	Phe	Lys	Asp	Ile	Asn	Asp	Val	Phe	Asp	Ala	Phe	Gln	Glu	Met
			180					185					190		
Ala	His	Asn	Val	Lys	Lys	Gly	Ile	Ile	Ala	Trp	Gly	Asp	Asp	Glu	His
		195					200					205			
Leu	Arg	Lys	Ile	Glu	Ala	Asp	Val	Pro	Ile	Tyr	Tyr	Tyr	Gly	Phe	Lys

210	215	220
Asp Ser Asp Asp Ile Tyr Ala Gln Asn Ile Gln Ile Thr Asp Lys Gly		
225	230	235 240
Thr Ala Phe Asp Val Tyr Val Asp Gly Glu Phe Tyr Asp His Phe Leu		
245	250	255
Ser Pro Gln Tyr Gly Asp His Thr Val Leu Asn Ala Leu Ala Val Ile		
260	265	270
Ala Ile Ser Tyr Leu Glu Lys Leu Asp Val Thr Asn Ile Lys Glu Ala		
275	280	285
Leu Glu Thr Phe Gly Gly Val Lys Arg Arg Phe Asn Glu Thr Thr Ile		
290	295	300
Ala Asn Gln Val Ile Val Asp Asp Tyr Ala His His Pro Arg Glu Ile		
305	310	315 320
Ser Ala Thr Ile Asp Thr Ala Arg Lys Lys Tyr Pro His Lys Glu Val		
325	330	335
Val Ala Val Phe Gln Pro His Thr Phe Ser Arg Thr Gln Ala Phe Leu		
340	345	350
Asn Glu Phe Ala Glu Ser Leu Cys Lys Ala Asp Arg Val Phe Leu Cys		
355	360	365
Glu Ile Phe Gly Ser Ile Arg Glu Asn Ser Gly Ala Leu Thr Ile Gln		
370	375	380
Asp Leu Ile Asp Lys Ile Gly Gly Ala Ser Phe Ile Asn Glu Asp Leu		
385	390	395 400
Ile Asn Val Leu Glu Gln Phe Asp Asn Ala Val Val Leu Phe Met Gly		
405	410	415
Ala Gly Asp Ile Gln Lys Leu Gln Asn Ala Tyr Leu Asp Lys Leu Gly		
420	425	430
Met Lys Asn Ala Phe		
435		

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 660 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATTTAAAGAT	TCGGATGACA	TTTATGCTCA	AATATTTCAA	ATTACGGATA	AAGGTACTGC	60
TGTTGATGTG	TATGTGGATG	GTGAGTTTTA	TGATCACTTC	CTGTCTCCAC	AATATGGTGA	120
CCATACAGTT	TTAAATGCAT	TAGCTGTAAT	TGCGATTAGT	TATTTAGAGA	AGCTAGATGT	180
TACAAATATT	AAAGAAGCAT	TAGAAACGTT	TGGTGGTGT	AAACGTCGTT	TCAATGAAAC	240
TACAATTGCA	AATCAAGTTA	TTGTAGATGA	TTATGCACAC	CATCCAAGAG	AAATTAGTGC	300
TACAATTGAC	ACAGCACGAA	AGAAATATCC	ACATAAAGAA	GTTGTTGCAG	TATTTCAACC	360
ACACACTTTC	TCTAGAACAC	AAGCATTTTT	AAATGAATTT	GCAGAAAGTT	TAAGTAAAGC	420
AGATCGTGTA	TTCTTATGTG	AAATTTTTTG	ATCAATTAGA	GAAAATACTG	GCGCATTAAC	480
GATACAAGAT	TTAATTGATA	AAATTGAAGG	TGCATCGTTA	ATTAATGAAG	ATTCTATTAA	540
TGTATTAGAA	CAATTGATA	ATGCTGTTGT	TTTATTTATG	GGTGCAGGTG	ATATTCAAAA	600
ATTACAAAAT	GCATATTTAG	ATAAATTAGG	CATGAAAAAT	GCGTTTTAAT	ATGTTTATAA	660

(2) INFORMATION FOR SEQ ID NO:4:

(A) LENGTH: 215 amino acids

(C) STRANDEDNESS: single

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Leu	Cys	Glu	Ile	Phe	Gly	Ser	Ile	Arg	Glu	Asn	Thr	Gly	Ala	Leu	Thr
145					150					155					160
Ile	Gln	Asp	Leu	Ile	Asp	Lys	Ile	Glu	Gly	Ala	Ser	Leu	Ile	Asn	Glu
				165					170					175	
Asp	Ser	Ile	Asn	Val	Leu	Glu	Gln	Phe	Asp	Asn	Ala	Val	Val	Leu	Phe
			180					185					190		
Met	Gly	Ala	Gly	Asp	Ile	Gln	Lys	Leu	Gln	Asn	Ala	Tyr	Leu	Asp	Lys
		195					200					205			
Leu	Gly	Met	Lys	Asn	Ala	Phe									
	210					215									

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CTTCATTAAT GAACGATGC

19

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GTTACAAATA TTAAAGAAG

19

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:

(i) an isolated polypeptide comprising an amino acid having at least:

- (a) 70% identity;
- (b) 80% identity;
- (c) 90% identity; or
- (d) 95% identity

to the amino acid sequence of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4;

(ii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2 or 4,

(iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2 or 4, and

(iv) a polypeptide which is encoded by a recombinant polynucleotide comprising the polynucleotide sequence of SEQ ID NO:1 or 3.

2. An isolated polynucleotide selected from the group consisting of:

(i) an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide that has at least

- (a) 70% identity;
- (b) 80% identity;
- (c) 90% identity; or
- (d) 95% identity;

to the amino acid sequence of SEQ ID NO:2 or 4, over the entire length of SEQ ID NO:2 or 4;

(ii) an isolated polynucleotide comprising a polynucleotide sequence that has at least:

- (a) 70% identity
- (b) 80% identity;
- (c) 90% identity; or
- (d) 95% identity;

over its entire length to a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 or 4;

(iii) an isolated polynucleotide comprising a nucleotide sequence which has at least:

- (a) 70% identity;
- (b) 80% identity;
- (c) 90% identity; or
- (d) 95% identity;

5 to that of SEQ ID NO: 1 or 3 over the entire length of SEQ ID NO:1 or 3;

(iv) an isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2 or 4;

(v) an isolated polynucleotide which is the polynucleotide of SEQ ID NO: 1 or 3;

10 (vi) an isolated polynucleotide obtainable by screening an appropriate library under stringent hybridization conditions with a probe having the sequence of SEQ ID NO: 1 or 3 or a fragment thereof;

(vii) an isolated polynucleotide encoding a mature polypeptide expressed by the MurC gene contained in the *Staphylococcus aureus*; and

15 (viii) a polynucleotide sequence complementary to said isolated polynucleotide of (i), (ii), (iii), (iv), (v), (vi) or (vii).

3. An antibody antigenic to or immunospecific for the polypeptide of claim 1.

4. A method for the treatment of an individual:

20 (i) in need of enhanced activity or expression of the polypeptide of claim 1 comprising the step of:

(a) administering to the individual a therapeutically effective amount of an agonist to said polypeptide; or

25 (b) providing to the individual an isolated polynucleotide comprising a polynucleotide sequence encoding said polypeptide in a form so as to effect production of said polypeptide activity *in vivo*; or

(ii) having need to inhibit activity or expression of the polypeptide of claim 1 comprising:

30 (a) administering to the individual a therapeutically effective amount of an antagonist to said polypeptide; or

(b) administering to the individual a nucleic acid molecule that inhibits the expression of a polynucleotide sequence encoding said polypeptide; or

- (c) administering to the individual a therapeutically effective amount of a polypeptide that competes with said polypeptide for its ligand, substrate, or receptor.

5 5. A process for diagnosing or prognosing a disease or a susceptibility to a disease in an individual related to expression or activity of the polypeptide of claim 1 in an individual comprising the step of:

(a) determining the presence or absence of a mutation in the nucleotide sequence encoding said polypeptide in the genome of said individual; or

10 (b) analyzing for the presence or amount of said polypeptide expression in a sample derived from said individual.

6. A method for screening to identify compounds that activate or that inhibit the function of the polypeptide of claim 1 which comprises a method selected from the group consisting of:

15 (a) measuring the binding of a candidate compound to the polypeptide or to the cells or membranes bearing the polypeptide or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;

(b) measuring the binding of a candidate compound to the polypeptide or to the cells or membranes bearing the polypeptide or a fusion protein thereof in the presence of a labeled competitor;

20 (c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes bearing the polypeptide;

(d) mixing a candidate compound with a solution containing a polypeptide of claim 1, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a standard;

(e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide and said polypeptide in cells, using for instance, an ELISA assay, or

25 (f) (1) contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second

component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound; and

(2) determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.

7. An agonist or an antagonist of the activity or expression polypeptide of claim 1.

8. An expression system comprising a polynucleotide capable of producing a polypeptide of claim

1 when said expression system is present in a compatible host cell.

9. A host cell comprising the expression system of claim 8 or a membrane thereof expressing a polypeptide selected from the group consisting of:

(i) an isolated polypeptide comprising an amino acid sequence selected from the group having at least:

- (a) 70% identity;
- (b) 80% identity;
- (c) 90% identity; or
- (d) 95% identity

to the amino acid sequence of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4;

(ii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2 or 4;

(iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2 or 4, and

(iv) a polypeptide which is encoded by a recombinant polynucleotide comprising the polynucleotide sequence of SEQ ID NO:1 or 3.

10. A process for producing a polypeptide selected from the group consisting of:

(i) an isolated polypeptide comprising an amino acid sequence selected from the group having at least:

- (a) 70% identity;
- (b) 80% identity;
- (c) 90% identity; or

(d) 95% identity

to the amino acid sequence of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4;

(ii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2 or 4;

5 (iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2 or 4, and

(iv) a polypeptide which is encoded by a recombinant polynucleotide comprising the polynucleotide sequence of SEQ ID NO:1 or 3,

comprising the step of culturing a host cell of claim 9 under conditions sufficient for the production of said polypeptide.

10

11. A process for producing a host cell comprising the expression system of claim 8 or a membrane thereof expressing a polypeptide selected from the group consisting of:

(i) an isolated polypeptide comprising an amino acid sequence selected from the group having at least:

15

(a) 70% identity;

(b) 80% identity;

(c) 90% identity; or

(d) 95% identity

to the amino acid sequence of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4;

20

(ii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2 or 4;

(iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2 or 4, and

(iv) a polypeptide which is encoded by a recombinant polynucleotide comprising the polynucleotide sequence of SEQ ID NO:1 or 3,

25

said process comprising the step of transforming or transfecting a cell with an expression system comprising a polynucleotide capable of producing said polypeptide of (i), (ii), (iii) or (iv) when said expression system is present in a compatible host cell such the host cell, under appropriate culture conditions, produces said polypeptide of (i), (ii), (iii) or (iv).

30

12. A host cell produced by the process of claim 11 or a membrane thereof expressing a polypeptide selected from the group consisting of:

(i) an isolated polypeptide comprising an amino acid sequence selected from the group having at least:

- (a) 70% identity;
- (b) 80% identity;
- (c) 90% identity; or
- (d) 95% identity

to the amino acid sequence of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4;

(ii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2 or 4;

(iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2 or 4, and
(iv) a polypeptide which is encoded by a recombinant polynucleotide comprising the polynucleotide sequence of SEQ ID NO:1 or 3.

13. A computer readable medium having stored thereon a member selected from the group consisting of: a polynucleotide comprising the sequence of SEQ ID NO. 1 or 3; a polypeptide comprising the sequence of SEQ ID NO. 2 or 4; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 1 or 3; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 2 or 4; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2 or 4; a polynucleotide comprising the sequence of SEQ ID NO. 1 or 3; a polypeptide comprising the sequence of SEQ ID NO. 2 or 4; a set of polynucleotide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 1 or 3; a set of polypeptide sequences wherein at least one of said sequences comprises the sequence of SEQ ID NO. 2 or 4; a data set representing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3; a data set representing a polynucleotide sequence encoding a polypeptide sequence comprising the sequence of SEQ ID NO. 2 or 4.

14. A computer based method for performing homology identification, said method comprising the steps of providing a polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3 in a computer readable medium; and comparing said polynucleotide sequence to at least one polynucleotide or polypeptide sequence to identify homology.

15. A further embodiment of the invention provides a computer based method for polynucleotide assembly, said method comprising the steps of: providing a first polynucleotide sequence comprising the sequence of SEQ ID NO. 1 or 3 in a computer readable medium; and
5 screening for at least one overlapping region between said first polynucleotide sequence and a second polynucleotide sequence.

16. An isolated polynucleotide selected from the group consisting of:

- 10 (a) an isolated polynucleotide comprising a nucleotide sequence which has at least 70%, 80%, 90%, 95%, 97% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;
- (b) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:3;
- (c) the polynucleotide of SEQ ID NO:3; or
- (d) an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide which has at least 70%, 80%, 90%, 95%, 97, 98, 99 or 99.5% identity to the amino acid sequence of SEQ ID
15 NO:4, over the entire length of SEQ ID NO:4.

17. A polypeptide selected from the group consisting of:


- (a) a polypeptide which comprises an amino acid sequence which has at least 70%, 80%, 90%, 95%, 97, 98, 99 or 99.5% identity to that of SEQ ID NO:4 over the entire length of SEQ ID
20 NO:4;
- (b) a polypeptide which has an amino acid sequence which is at least 70%, 80%, 90%, 95%, 97, 98, 99 or 99.5% identity to the amino acid sequence of SEQ ID NO:4 over the entire length of SEQ ID NO:4;
- (c) a polypeptide which comprises the amino acid of SEQ ID NO:4;
- 25 (d) a polypeptide which is the polypeptide of SEQ ID NO:4;
- (e) a polypeptide which is encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:3.

[illegible]

- 73 -

United States Patent & Trademark Office

Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

1. Application papers are not suitable for scanning and are not in compliance with 37 CFR 1.52 because:
 - ☐ All sheets must be the same size and either A4 (21 cm x 29.7 cm) or 8-1/2" x 11". Pages _____ do not meet these requirements.
 - ☐ Papers are not flexible, strong, smooth, non-shiny, durable, and white.
 - ☐ Papers are not typewritten or mechanically printed in permanent ink on one side.
 - ☐ Papers contain improper margins. Each sheet must have a left margin of at least 2.5 cm (1") and top, bottom and right margins of at least 2.0 cm (3/4").
 - ☐ Papers contain hand lettering.
2. Drawings are not in compliance and were not scanned because:
 - ☐ The drawings or copy of drawings are not suitable for electronic reproduction.
 - ☐ All drawings sheets are not the same size. Pages must be either A4 (21 cm x 29.7 cm) or 8-1/2" x 11".
 - ☐ Each sheet must include a top and left margin of at least 2.5 cm (1"); a right margin of at least 1.5 cm (9/16") and a bottom margin of at least 1.0 cm (3/8").
3. Page(s) _____ are not of sufficient clarity, contrast and quality for electronic reproduction.
4. Page(s) _____ are missing.
5. OTHER: No Declarations/No Drawings